

TORELLI'S THEOREM FOR HIGH DEGREE SYMMETRIC PRODUCTS OF CURVES

NAJMUDDIN FAKHRUDDIN

ABSTRACT. We show that two smooth projective curves C_1 and C_2 of genus g which have isomorphic symmetric products are isomorphic unless $g = 2$. This extends a theorem of Martens.

Let k be an algebraically closed field and C_1 and C_2 two smooth projective curves of genus $g > 1$ over k . It is a consequence of Torelli's theorem that if $Sym^{g-1}C_1 \cong Sym^{g-1}C_2$, then $C_1 \cong C_2$. The same holds for the d 'th symmetric products, for $1 \leq d < g - 1$ as a consequence of a theorem of Martens [2]. In this note we shall show that with one exception the same result continues to hold for all $d \geq 1$, i.e. we have the following

Theorem 1. *Let C_1 and C_2 be smooth projective curves of genus $g \geq 2$ over an algebraically closed field k . If $Sym^d C_1 \cong Sym^d C_2$ for some $d \geq 1$, then $C_1 \cong C_2$ unless $g = d = 2$.*

It is well known that there exist non-isomorphic curves of genus 2 over $\mathbf{C}$ with isomorphic Jacobians. Since the second symmetric power of a genus 2 curve is isomorphic to the blow up of the Jacobian in a point, it follows that our result is the best possible.

Proof of Theorem. Let C_i , $i = 1, 2$ be two curves of genus $g > 1$ with $Sym^d C_1 \cong Sym^d C_2$ for some $d \geq 1$. Since the Albanese variety of $Sym^d C_i$, $d \geq 1$, is isomorphic to the Jacobian $J(C_i)$, it follows that $J(C_1) \cong J(C_2)$. If $d \leq g - 1$, the theorem follows immediately from [2], since the image of $Sym^d C_i$ in $J(C_i)$ (after choosing a basepoint) is $W_d(C_i)$. Note that in this case it suffices to have a birational isomorphism from $Sym^d C_1$ to $Sym^d C_2$.

Suppose $g \leq d \leq 2g - 3$. Then the Albanese map from $Sym^d C_i$ to $J(C_i)$ is surjective with general fibre of dimension $d - g$. Interpreting the fibres as complete linear systems of degree d on C_i , it follows by Serre duality that the subvariety of $J(C_i)$ over which the fibres are of dimension $> d - g$ is isomorphic to $W_{2g-2-d}(C_i)$. Therefore if $Sym^d C_1 \cong Sym^d C_2$, then $W_{2g-2-d}(C_1) \cong W_{2g-2-d}(C_2)$, so Martens' theorem implies that $C_1 \cong C_2$.

Now suppose that $d > 2g - 2$ and $g > 2$. By choosing some isomorphism we identify $J(C_1)$ and $J(C_2)$ with a fixed abelian variety A . If $\phi : Sym^d C_1 \rightarrow Sym^d C_2$ is our given isomorphism, from the universal property of the Albanese morphism we obtain a commutative diagram

$$\begin{array}{ccc} Sym^d C_1 & \xrightarrow{\phi} & Sym^d C_2 \\ \pi_1 \downarrow & & \downarrow \pi_2 \\ A & \xrightarrow{f} & A \end{array}$$

1

where the π_i 's are the Albanese morphisms corresponding to some basepoints and f is an automorphism of A (not necessarily preserving the origin). By replacing C_2 with $f^{-1}(C_2)$ we may then assume that f is the identity.

Since $d > 2g - 2$, the maps π_i , $i = 1, 2$ make $Sym^d C_i$ into projective bundles over A . By a theorem of Schwarzenberger [5], $Sym^d C_i \cong Proj(E_i)$, where E_i is a vector bundle on A of rank $d - g + 1$ with $c_j(E_i) = [W_{g-j}(C_i)]$, $i = 1, 2$, $0 \leq j \leq g - 1$, in the group of cycles on A modulo numerical equivalence. Since ϕ is an isomorphism of projective bundles, it follows that there exists a line bundle L on A such that $E_1 \cong E_2 \otimes L$.

Let $\theta_i = [W_{g-1}(C_i)]$, so by Poincaré's formula $[W_{g-j}(C_i)] = \theta_i^j / j!$, $i = 1, 2$, $i \leq j \leq g - 1$. The lemma below implies that $\theta_i^{g-1} = \theta_2^{g-1}$ in the group of cycles modulo numerical equivalence on A . Since $\theta_i^g = g!$, this implies that $\theta_1 \cdot [C_2] = g$. By Matsusaka's criterion [3], it follows that $W_{g-1}(C_1)$ is a theta divisor for C_2 , which by Torelli's theorem implies that $C_1 \cong C_2$.

If $d = 2g - 2$ and $g > 2$, then we can still apply the previous argument. In this case we also have that $Sym^d(C_i) \cong Proj(E_i)$, $i = 1, 2$ but E_i is a coherent sheaf which is not locally free. However on the complement of some point of A it does become locally free and the previous formula for the Chern classes remains valid.

The above argument clearly does not suffice if $g = 2$. To handle this case we shall use some properties of Picard bundles for which we refer the reader to [4]. Suppose that $d > 2$ and C_i , $i = 1, 2$ are two *non-isomorphic* curves of genus 2 with $Sym^d C_1 \cong Sym^d C_2$. Using the same argument (and notation) as the $g > 2$ case, it follows that there exist embeddings of C_i , $i = 1, 2$, in A and a line bundle L on A such that $E_1 \cong E_2 \otimes L$ and $L^{\otimes d-1} \cong \mathcal{O}(C_1 - C_2)$ (we identify C_i , $i = 1, 2$ with their images).

For $i \geq 1$, let G_i denote the i -th Picard sheaf associated to C_2 , so that $Proj(G_i) \cong Sym^i(C_2)$. (G_i is the sheaf denoted by F_{2-i} in [4] and $G_d \cong E_2$). There is an exact sequence ([4, p.172]):

$$(1) \quad 0 \rightarrow \mathcal{O}_A \rightarrow G_i \rightarrow G_{i-1} \rightarrow 0$$

for all $i > 1$. We will use this exact sequence and induction on i to compute the cohomology of sheaves of the form $E_1 \otimes P \cong E_2 \otimes L \otimes P$, where $P \in Pic^0(A)$.

Consider first the cohomology of G_1 , which is the pushforward of a line bundle of degree 1 on a translate of C_2 . Since we have assumed that $C_1 \not\cong C_2$, it follows that $C_1 \cdot C_2 > 2$. Since $C_1^2 = C_2^2 = 2$, $deg(L|_{C_2}) = (C_1 - C_2) \cdot C_2 / (d - 1) > 0$. By Riemann-Roch it follows that $h^j(A, G_1 \otimes L \otimes P)$, $j = 1, 2$ is independent of P , except possibly for one P if $deg(L|_{C_2}) = 1$, and $h^2(A, G_1 \otimes L \otimes P) = 0$ since G_1 is supported on a curve.

Now $C_1 \cdot C_2 > 2$ also implies that $c_1(L)^2 < 0$. By the index theorem, it follows that $h^0(A, L \otimes P) = h^2(A, L \otimes P) = 0$ and $h^1(A, L \otimes P)$ is independent of P . Therefore by tensoring the exact sequence (1) with $L \otimes P$ and considering the long exact sequence of cohomology, we obtain an exact sequence

$$(2) \quad \begin{aligned} 0 \rightarrow H^0(A, G_i \otimes L \otimes P) &\rightarrow H^0(A, G_{i-1} \otimes L \otimes P) \rightarrow H^1(A, L \otimes P) \\ &\rightarrow H^1(A, G_i \otimes L \otimes P) \rightarrow H^1(A, G_{i-1} \otimes L \otimes P) \rightarrow 0 \end{aligned}$$

and isomorphisms $H^2(A, G_i \otimes L \otimes P) \rightarrow H^2(A, G_{i-1} \otimes L \otimes P)$ for all $i > 1$. By induction, it follows that $H^2(A, G_i \otimes L \otimes P) = 0$ for all $i > 0$. Since the Euler characteristic of $G_i \otimes L \otimes P$ is independent of P , the above exact sequence (2)

along with induction shows that for all $i > 0$ and $j = 0, 1, 2$, $h^j(G_i \otimes L \otimes P)$ is independent of P , except for possibly one P . In particular, this holds for $i = d$ hence $h^j(A, E_1 \otimes P)$ is independent of P except again for possibly one P . We obtain a contradiction by using the computation of the cohomology of Picard sheaves in Proposition 4.4 of [4]: This implies that $h^1(A, E_1 \otimes P)$ is one or zero depending on whether the point in A corresponding to P does or does not lie on a certain curve (which is a translate of C_1).

□

Lemma 1. *Let X be an algebraic variety of dimension $g \geq 3$ and let E_i , $i = 1, 2$ be vector bundles on X of rank r . Suppose $c_1(E_i) = \theta_i$, $c_j(E_i) = \theta_i^j/j!$ for $i = 1, 2$ and $j = 2, 3$ ($j = 2$ if $g = 3$), and $E_1 \cong E_2 \otimes L$ for some line bundle L on X . Then $\theta_1^j = \theta_2^j$ for all $j > 1$ ($j = 2$ if $g = 3$).*

Proof. Since $E_1 \cong E_2 \otimes L$, $c_1(E_1) = c_1(E_2) + rc_1(L)$, hence $c_1(L) = (\theta_1 - \theta_2)/r$.

For a vector bundle E of rank r and a line bundle L on any variety, we have the following formula for the Chern polynomial ([1], page 55):

$$c_t(E \otimes L) = \sum_{k=0}^r t^k c_t(L)^{r-k} c_i(E).$$

Letting $E = E_1$, $E \otimes L = E_2$, and expanding out the terms of degree 2 and 3, one easily sees that $\theta_1^j = \theta_2^j$ for $j = 2$ and also for $j = 3$ if $g > 3$. (Note that this only requires knowledge of $c_j(E_i)$ for $j = 1, 2, 3$). Since any integer $n > 1$ can be written as $n = 2a + 3b$ with $a, b \in \mathbf{N}$, the lemma follows. □

We do not know whether the theorem holds if k is not algebraically closed.

Remark. S. Ramanan has suggested it may also be possible to prove the theorem when $g > 2$ by computing the Chern classes of the push forward of the tangent bundle of $\text{Sym}^d C$ to $J(C)$.

Acknowledgements. We thank A. Collino for some helpful correspondence, in particular for informing us about the paper [2], and S. Ramanan for the above remark.

REFERENCES

- [1] W. FULTON, *Intersection theory*, Springer-Verlag, Berlin, second ed., 1998.
- [2] H. H. MARTENS, *An extended Torelli theorem*, Amer. J. Math., 87 (1965), pp. 257–261.
- [3] T. MATSUSAKA, *On a characterization of a Jacobian variety*, Memo. Coll. Sci. Univ. Kyoto. Ser. A. Math., 32 (1959), pp. 1–19.
- [4] S. MUKAI, *Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves*, Nagoya Math. J., 81 (1981), pp. 153–175.
- [5] R. L. E. SCHWARZENBERGER, *Jacobians and symmetric products*, Illinois J. Math., 7 (1963), pp. 257–268.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

E-mail address: naf@math.tifr.res.in